

Appl. No. : 09/859,651
Filed : May 17, 2001

42. (Previously added) The method of claim 41, wherein the protein is human alpha-2b.
43. (Previously added) The method of claim 41, wherein the host cell further comprises recA¹³.
44. (Previously added) The method of claim 41, wherein the *E. coli* host cell produces a suppressor for the repair of amber-mutations.
45. (Previously added) The method of claim 41, wherein the *E. coli* host cell lacks a suppressor for the repair of amber-mutations.
46. (Previously added) The method of claim 41, wherein the infecting bacteriophage λ is provided at a multiplicity of infection in a range of about 1 to about 100.
47. (Previously added) The method of claim 41, wherein the infecting bacteriophage λ is provided at a multiplicity of infection in a range of about 10 to about 25.
48. (Cancel)
49. (Currently amended) A method for producing a soluble biologically-active protein, comprising:
- transforming an *E. coli* host cell with a plasmid having at least one copy of an expressible eukaryotic gene encoding said protein;
 - lytically infecting the transformed bacterial host cell with a bacteriophage λ having cI₈₅₇, Q_{am117}, and R_{am54} mutations, wherein the bacteriophage also contains at least one copy of said expressible gene encoding said protein; and
 - cultivating the *E. coli* host cell under a culture condition that induces lytic growth of said cell without lysis to produce the soluble, biologically-active protein.
50. (Previously added) The method of claim 49, wherein the strain of *E. coli* produces a suppressor for repairing amber-mutations.

Appl. No. : 09/859,651
Filed : May 17, 2001

51. (Previously added) The method of claim 49, wherein the strain of *E. coli* lacks a suppressor for repairing amber-mutations.

52. (Previously added) The method of claim 49, wherein said protein is human alpha-2b interferon.

53. (Currently amended) A method for producing a biologically active protein, comprising:

transforming an *E. coli* host cell with a plasmid having at least one copy of an expressible eukaryotic gene encoding said protein;

lytically infecting the transformed bacterial host cell with a bacteriophage λ having at least one mutated gene selected from the group consisting of N, Q, and R; and

cultivating the *E. coli* host cell under a culture condition that induces lytic growth of said cell without lysis to produce the biologically active protein.

54. (Previously added) The method of claim 53, wherein the bacteriophage λ has a temperature-sensitive mutation.

55. (Previously added) The method of claim 54, wherein the temperature-sensitive mutation is *cl*₈₅₇.

56. (Previously added) The method of Claim 53, wherein said strain of *E. coli* lacks a suppressor for repairing amber-mutations.

57. (Previously added) The method of Claim 53, wherein said strain of *E. coli* is *recA* deficient.

58. (Currently amended) A method for producing a biologically active protein, comprising:

transforming an *E. coli* host cell with a plasmid having at least one copy of an expressible eukaryotic gene encoding said protein;

lytically infecting the transformed bacterial host cell with a bacteriophage λ , having at least one mutated gene selected from the group consisting of N, Q, and R,

wherein the bacteriophage also contains at least one copy of said expressible gene encoding said protein; and

SUB c1 Cont
cultivating the *E. coli* host cell under a culture condition that induces lytic growth of said cell without lysis until production of said protein is reached.

59. (Previously added) The method of claim 58, wherein the bacteriophage λ has a temperature-sensitive mutation.

60. (Previously added) The method of claim 59, wherein the temperature-sensitive mutation is *cl₈₅₇*.

By Care
61. (Previously added) The method of Claim 58, wherein said *E. coli* host cell lacks a suppressor for repairing amber-mutations.

62. (Previously added) The method of Claim 58, wherein said *E. coli* host cell is *recA* deficient.

63. (Currently amended) A method of producing a biologically-active protein comprising:
growing a first strain of *E. coli* cells, which harbor a strain of bacteriophage λ ,
wherein the bacteriophage λ has a temperature-sensitive mutation,
manipulating the temperature to provide for lysis of the first strain of *E. coli* cells and release of the bacteriophage λ ,

SUB c1
adding the released bacteriophage λ to a second strain of *E. coli* cells to lytically infect the second strain of *E. coli* cells with the released bacteriophage λ , wherein said second strain of *E. coli* cells has been transformed with a plasmid having at least one copy of an expressible gene encoding said biologically-active protein; and

culturing the second strain of *E. coli* host cells such that said biologically-active protein is produced and released to the media as a soluble, biologically-active protein.

64. (Previously added) The method of claim 63, wherein the temperature-sensitive mutation is *cl₈₅₇*.